



Factors Affecting Interpretation of Postprandial Glucose and Insulin Areas

MARY C. GANNON, PhD, AND FRANK Q. NUTTALL, MD, PhD

Recently there has been an increased interest in determining the circulating glucose concentration after the ingestion of various individual foods and mixed meals. The purpose of these determinations is to systematically rank foods with respect to their quantitative effect on postmeal glucose concentration. Potentially such data could be useful in designing a diet for individuals with diabetes. We believe this concept is good. However, several factors that may affect interpretation of the data used to develop this ranking need to be considered before the utility of this approach to dietary management can be assessed: 1) duration of time over which the data are collected and analyzed; 2) use of absolute versus incremental areas in the determinations; 3) inclusion or exclusion of negative areas if incremental areas are used; 4) differences in response to a given food in males compared with females; 5) severity of diabetes; 6) confounding effects of oral agents or insulin treatment; 7) reproducibility of data; 8) differences in collection of blood sample; 9) food composition, processing, and preparation; 10) the dose-response relationship to ingestion of a given carbohydrate; 11) the meal being studied, i.e., first, second, or third meal of the day; and 12) a possible effect of the composition of the previous meal, if the response is tested to any meal other than the first meal of the day. *Diabetes Care* 10:759-63, 1987

For the past several years there has been an interest in determining the circulating glucose concentrations in peripheral blood after the ingestion of individual foods (1-5). These studies have been conducted in normal and diabetic subjects. The data obtained from such studies have been analyzed by quantitating the glucose response to a known amount of carbohydrate in a particular food. A common way of calculating this has been to determine a glycemic index. The glycemic index has been defined by Jenkins et al. (3) as the area under the glucose curve after a 50-g carbohydrate test meal divided by the area under the glucose curve after a meal consisting of a standard reference food containing 50 g carbohydrate, multiplied by 100. Originally, blood glucose area was determined over a 2-h period, and 50 g glucose was used as the standard reference meal (3). Recently, blood glucose determinations of diabetic subjects have been performed over a 3-h period, and the standard reference meal has been 50 g carbohydrate in the form of bread (6). Such data have been used to assign a relative ordering of plasma glucose responses to various carbohydrate-containing foods. The ultimate purpose of this ordering is to assist the diabetic individual in meal planning.

Theoretically, ingestion of foods of low glycemic index could be useful in the dietary management of diabetes.

Recently, the validity and clinical utility of this index has been called into question. When foods of varying glycemic index were incorporated into a mixed meal, the glycemic index was not useful in predicting the glucose response (7-10). However, this has not been a universal finding (11-14). In our laboratory, we have been interested in quantitating both the insulin and glucose responses to foods as accurately as possible. During these studies, we became aware of a potential source of error in determining the relative glucose area (RGA) after the ingestion of various foods. This potential source of error is the time duration over which the plasma glucose area is determined after a meal. Subsequent review of other studies has indicated several potentially confounding variables that should be considered in interpretation of glycemic index data.

In this report, we use the term RGA instead of *glycemic index*, because the time over which relative areas are determined is variable. RGA is defined as the net glucose area after the ingestion of a test meal, divided by the net glucose area after the ingestion of 50 g glucose, multiplied by 100.

This may be determined at any time point after the ingestion of a single food or a mixed meal. The fasting glucose concentration is used as a baseline. The glycemic index was originally based on 2 h of data with glucose as the reference food (3) and now is based on 2 h of data in normal subjects or 3 h of data in diabetic subjects with bread as the reference food (15).

When we began our studies of glucose and insulin responses to food ingestion, we determined the time required for peripheral plasma glucose and peripheral serum insulin concentrations to return to premeal values after the ingestion of 50 g glucose. We observed that the time required for the glucose concentrations to return to the fasting baseline concentration was ~1.5 h in normal individuals (16). Insulin concentrations returned to baseline at ~3 h. In untreated mildly non-insulin-dependent diabetic (NIDDM) male subjects, ~4–5 h was required for plasma glucose to return to the fasting concentration (17,18). Insulin concentrations were still modestly elevated 5 h after ingestion of 50 g glucose in these subjects. However, we did not consider it practical to continue the studies for a longer period.

Knowing that the rate of digestion and absorption of foods may be different, it was necessary to arbitrarily choose a time

frame during which to quantitate the glucose and insulin responses to various other foods. We chose to collect and analyze data for 5 h after ingestion of the test foods in the diabetic patients, because 5 h generally was required for the glucose concentration to return to the fasting level after 50 g glucose, the standard reference meal in our study. Five hours appears to be the longest period monitored in any published study. We decided it would be useful to determine the effect on RGA when the data were quantitated over various periods up to 5 h.

It was apparent that, at least for some foods, RGA was not constant with time (Fig. 1), and the differences in RGA often were quite large. For example, after the ingestion of sucrose, the RGAs at 1, 3, and 5 h were 116, 67, and 44, respectively (Fig. 1A); i.e., when the glucose area after the ingestion of sucrose was determined at 1 h and compared with the glucose area after the ingestion of glucose at 1 h, the postsucrose area was 116% that of glucose. Decreases in RGA with time also were observed after fructose and milk ingestion (Fig. 1A), and similar results were obtained when the data for apples, apple juice, and orange juice were analyzed (data not shown). Thus, sucrose-, fructose-, and lactose-containing foods generally resulted in a decrease in RGA




FIG. 1. Relative glucose area (RGA) after ingestion of various foods. A: decreasing RGA with time after ingestion of 50 g CHO in form of sucrose, fructose, or milk in untreated non-insulin-dependent diabetic (NIDDM) male subjects ($n = 7$) (17). B: increasing RGA with time after ingestion of 50 g CHO in form of lentils or kidney beans in untreated NIDDM male subjects ($n = 7$) (18). C: little change in RGA with time after ingestion of 50 g CHO in form of potatoes, bread, oatmeal, or rice in untreated NIDDM male subjects ($n = 7$) (18). D: decreasing RGA with time after ingestion of glucose plus protein meal. Normal subjects ($n = 8$) ingested 50 g glucose plus 50 g protein (lean beef) (16); diabetic subjects ($n = 6$) ingested 50 g glucose plus 25 g protein (cottage cheese) (35).

with time. On the other hand, with slowly digested foods such as legumes and kidney beans, RGA increased with time (Fig. 1B; 18).

Ingestion of some foods resulted in little change in the RGA with time, because the peak response and time required for plasma glucose to return to the fasting concentration was similar to that after ingestion of pure glucose (Fig. 1C; 18). Protein ingestion does not increase plasma glucose concentration but stimulates insulin secretion (19,20). Thus, the additional insulin secreted in response to protein when given with glucose could influence RGA. Indeed, this was the case. In both normal and NIDDM individuals, RGA after ingestion of glucose and protein decreased with time (Fig. 1D). This primarily was due to a more rapid return of the glucose concentration to baseline when 50 g protein in the form of lean beef was ingested with 50 g glucose (16). Thus, the protein in a mixed meal is likely to influence RGA, and the effect would be more prominent at later time points.

We and others have demonstrated little effect of dietary fat on the plasma glucose response after the first meal of the day in normal subjects (21) and insulin-dependent diabetic patients (22). However, the 1-h postprandial glucose area was significantly less after the ingestion of 50 g fat given with 50 g carbohydrate compared with 50 g carbohydrate alone in normal subjects (23). In NIDDM patients, the presence of fat in the meal modestly decreased the postprandial glucose area as well (22). Furthermore, the presence of a large amount of fat in the meal resulted in a delay of the glucose peak of the second and third meals of the day (21).

Theoretically, all studies should be conducted until RGA remains essentially unchanged with time. However, conducting studies over the time necessary to achieve this may be impractical. A compromise may be to select a time at which the rate of change approaches a minimum. Alternatively, the glucose response to various foods may be studied with the usual duration of time between meals in that population. In our society, meals are generally ingested 4–6 h apart.

Diabetes presents a particular problem because it affects the duration of time over which the postmeal glucose concentration remains elevated, even though it does not significantly influence the rate of digestion or absorption of foods. Both the duration and magnitude of elevation may be influenced by the severity of the insulin insufficiency. Theoretically, the severity of diabetes also could affect the metabolism of fructose and galactose. Therefore, the response to various foods should be standardized in the study population. In addition, it may be advisable to test the effect of ingestion of pure monosaccharides and foods containing those monosaccharides on the study population (17). In diabetic patients receiving insulin or oral hypoglycemic agents, the effect of the medication may vary from day to day (24), which also could complicate interpretation of the data.

Other factors should be considered in interpreting RGA data. The area under the glucose curve has been determined with the absolute values obtained for plasma glucose (2,25) or by determining the incremental glucose area above the

fasting concentration (3,6,11–13,15–18). As pointed out by Wolever and Jenkins (11), any differences between responses to foods will appear smaller if the absolute areas are determined. In addition, differences in fasting glucose values may strongly influence absolute areas but have little effect on incremental area unless the glucose exceeds the renal threshold for glucose excretion. An elevated glucose concentration should stimulate more non-insulin-dependent glucose uptake by muscle (26). Theoretically, this could affect RGA. Whether this is quantitatively important has not been determined, but it probably is not of major importance except in poorly controlled diabetic subjects.

When incremental areas are used, inclusion or exclusion of negative values also may have a considerable effect on RGA of a particular food. With incremental areas, we have observed values below baseline on several occasions, particularly after foods containing fructose or milk. Thus, the method of analysis should be clearly stated.

When the blood glucose response is determined after the ingestion of oral glucose loads given ~4 h apart to normal subjects, the response to the second meal is smaller than to the first. When glucose loads are given in succession, the facilitated disposal of the second and subsequent glucose loads is the so-called Staub-Traugott effect (27–29). Thus, the time of day and the meal sequence may influence the response. In most studies the subjects are given the test meal for breakfast. However, in one study a mixed meal was given for lunch, and the composition of the breakfast was not stated (2). Amplification of a glucose second-meal effect also may occur in normal subjects and NIDDM patients when protein is included in the meal (30).

The glycemic index for single foods has been determined based on an amount of food that contains a defined quantity of carbohydrate, usually 50 g. Whether the glucose response to ingestion of a particular food is linearly related to the amount of food ingested (and thus the quantity of carbo-




FIG. 2. Relative glucose area after ingestion of various amounts of glucose in untreated non-insulin-dependent diabetic male subjects ($n = 7$). □, 15 g glucose; ◆, 25 g glucose; ■, 35 g glucose.

hydrate) is another important consideration. This is particularly relevant when applying the glycemic index to mixed meals in which the amount of a food ingested is less than that used to determine the glycemic index. As indicated in Fig. 2, RGA is not constant with time when 15, 25, and 35 g glucose were ingested by untreated NIDDM patients. In addition, the slope of RGA with time was not constant from one dose to another, indicating that as a greater quantity of carbohydrate is ingested, plasma glucose concentration tends to remain elevated longer. These confounding factors should be considered when comparing data from various laboratories and mixed-meal responses with those anticipated from the glycemic index of various carbohydrate-containing foods in the mixed meal.

The variability of an individual's response to replicate testing also must be considered. We have very little information on this subject, but our available data from NIDDM patients indicate that the standard error can be quite large. We have no information on replicate testing of other foods. Others reported that when five normal subjects ingested two rice meals ~2 yr apart, glucose areas differed by 0–23% (15).

Use of capillary versus venous blood sampling also may result in problems in comparing data. Capillary blood is used directly for glucose determination with enzyme-impregnated plastic strips. Venous blood is centrifuged, and the serum or plasma is used for enzymatic glucose determination. Both assay methods are specific for glucose, but the absolute glucose concentration measured may differ. Although glucose is freely diffusible between plasma and erythrocytes and the glucose concentration is the same in both plasma water and erythrocyte water, the water content of the erythrocyte is ~72%, whereas that of plasma is ~94% (31). Thus, the true glucose concentration in whole blood is a function of the hematocrit. For 100 mg/dl plasma glucose, whole-blood glucose would be 89 mg/dl based on a hematocrit of 45; i.e., plasma would be higher by 11%. On the other hand, the glucose concentration may be lower in venous blood drawn from an antecubital vein compared with capillary blood from a finger because of removal of glucose by skin and skeletal muscle in the forearm.

We have determined another potential problem in interpreting the glucose response to foods. The glucose response to a mixed meal was different in normal men and women (21). In women the plasma glucose was transiently elevated for 1 h after breakfast. Over the subsequent 3 h, glucose concentration was below the fasting baseline. This biphasic response was not present in normal men. Consequently, 4-h glucose area after a mixed meal was $-10 \text{ mg} \cdot \text{h} \cdot \text{dl}^{-1}$ in women but $+70 \text{ mg} \cdot \text{h} \cdot \text{dl}^{-1}$ in men. Whether this sex difference exists in diabetic individuals has not been determined. Also, we are not aware of studies designed to determine a potential sex difference in the glucose response to individual foods.

For starch-containing foods, food processing and preparation may affect RGA. For example, if starch is extracted from beans, it is rapidly digested, whereas in ordinary cooked beans, digestion is slow and incomplete (32). When 50–75

g of raw starch was administered by mouth to normal subjects, no change in blood glucose was observed. However, when cooked starch was administered, blood glucose clearly increased (33). Starch exists in plants in the form of granules, the structure of which is characteristic for each plant species. Before starch can be hydrolyzed in vivo, the granule must be disrupted. This generally is done by heating in water, which causes the granules to swell, resulting in disruption of the structure. However, the ease with which disruption is accomplished varies considerably. Generally, starches with a high amylose-to-amylopectin ratio are more rigid and resistant to disruption than are starches with a low amylose-to-amylopectin ratio. For example, ingestion of a 50-g carbohydrate meal of muffins made from low-amylose cornstarch results in an increase in plasma glucose that is even greater than with 50 g pure glucose (117%), whereas ingestion of a 50-g carbohydrate meal of muffins made from high-amylose cornstarch was only 62% of that obtained with 50 g pure glucose (19). The only difference in the two meals was the type of cornstarch used to make the muffins. This clearly demonstrates that starch structure has effects on the postmeal plasma glucose response. In fruits such as bananas, the degree of ripeness may be a significant factor in the response (34).

In summary, there are numerous factors that may influence RGA and thus the interpretation of glycemic-index data. The concept of systematically ranking foods with respect to the quantitative effect on blood glucose is good and should prove useful in designing a diet for diabetic individuals and in educating patients. However, problems that may affect interpretation of such data must be recognized. These fall into three broad categories: the method of determining the glucose area, the patient population and method of sample collection, and the foods tested. From the number of factors identified that may influence the interpretation of postprandial glucose and insulin areas, additional, carefully designed research is necessary before single-food RGA data may be logically applied to meal planning for individuals with diabetes.

ACKNOWLEDGMENTS: This study was supported by a grant from the Veterans Administration and funds from the National Dairy Board, administered in cooperation with the National Dairy Council.

From the Metabolic-Endocrine Section, Veterans Administration Medical Center, and the Department of Medicine, University of Minnesota, Minneapolis, Minnesota.

Address correspondence and reprint requests to Mary C. Gannon, PhD, Nutritional Biochemist, Endocrine Research (111G), V.A. Medical Center, Minneapolis, MN 55417.

REFERENCES

1. Crapo PA, Reaven G, Olefsky J: Plasma glucose and insulin responses to orally administered simple and complex carbohydrates. *Diabetes* 25:741–47, 1976
2. Coulston A, Greenfield M, Kraemer F, Tobey T, Reaven G: Effect of source of dietary carbohydrate on plasma glucose and insulin responses to test meals in normal subjects. *Am J Clin Nutr* 33:1279–82, 1980

3. Jenkins DJA, Wolever TMS, Taylor RH, Barker H, Fielden H, Baldwin JM, Bowling AC, Newman HC, Jenkins AL, Goff DV: Glycemic index of foods: a physiological basis for carbohydrate exchange. *Am J Clin Nutr* 34:362-66, 1981
4. Nuttall FQ, Mooradian AD, DeMarais R, Parker S: The glycemic effect of different meals approximately isocaloric and similar in protein, carbohydrate, and fat content as calculated using the ADA exchange lists. *Diabetes Care* 6:432-35, 1983
5. Bantle JP, Laine DC, Castle GW, Thomas W, Hoogwerf BJ, Goetz FC: Postprandial glucose and insulin responses to meals containing different carbohydrates in normal and diabetic subjects. *N Engl J Med* 309:7-12, 1983
6. Jenkins DJA, Wolever TMS, Jenkins AL, Thorne MJ, Lee R, Kalmusky J, Reichert R, Wong GS: The glycaemic index of foods tested in diabetic patients: a new basis for carbohydrate exchange favoring the use of legumes. *Diabetologia* 24:257-64, 1983
7. Coulston AM, Hollenbeck CB, Reaven M: Utility of studies measuring glucose and insulin responses to various carbohydrate-containing foods. *Am J Clin Nutr* 39:163-65, 1984
8. Coulston AM, Hollenbeck CB, Liu GC, Williams RA, Starich GH, Mazzaferri EL, Reaven GM: Effect of source of dietary carbohydrate on plasma glucose, insulin, and gastric inhibitory polypeptide responses to test meals in subjects with non-insulin-dependent diabetes mellitus. *Am J Clin Nutr* 40:965-70, 1984
9. Coulston A, Hollenbeck C: Comparison of plasma glucose and insulin responses to mixed meals of predicted high, medium, and low glycemic response (Abstract). *Diabetes* 35 (Suppl. 1):43A, 1986
10. Laine DC, Thomas JW, Bantle JP: Comparison of the predictive capabilities of the diabetic exchange lists and the glycemic indices of foods (Abstract). *Diabetes* 35 (Suppl. 1):43A, 1986
11. Wolever TMS, Jenkins DJA: The use of the glycemic index in predicting the blood glucose response to mixed meals. *Am J Clin Nutr* 43:167-72, 1986
12. Wolever TMS, Collier GR, Wong GS, Josse RG: Prediction of glycemic response to mixed meals in NIDDM (Abstract). *Diabetes* 35 (Suppl. 1):43A, 1986
13. Collier GR, Wolever TMS, Wong GS, Josse RG: Prediction of glycemic response to mixed meals in non-insulin-dependent diabetic subjects. *Am J Clin Nutr* 44:349-52, 1986
14. Slama G, Bornet F, Blayo A, Costagliola D, Haardt MJ, Tchobroutsky G: Insulinogenic and glycaemic indexes of various starch-rich foods taken in a mixed meal or alone in type 2 diabetics (Abstract). *Diabetes* 34 (Suppl. 1):48A, 1985
15. Wolever TMS, Nuttall FQ, Lee R, Wong GS, Josse RG, Csima A, Jenkins DJA: Prediction of the relative blood glucose response to mixed meals using the white bread glycemic index. *Diabetes Care* 8:418-28, 1985
16. Krezowski PA, Nuttall FQ, Gannon MC, Bartosh NH: The effect of protein ingestion on the metabolic response to oral glucose in normal individuals. *Am J Clin Nutr* 44:847-56, 1986
17. Gannon MC, Nuttall FQ, Krezowski PA, Billington CJ, Parker S: The serum insulin and plasma glucose responses to milk and fruit products in type 2 diabetic patients. *Diabetologia* 29:784-91, 1986
18. Krezowski PA, Nuttall FQ, Gannon MC, Billington CJ, Parker S: Insulin and glucose responses to various starch-containing foods in type II diabetic subjects. *Diabetes Care* 10:205-12, 1987
19. Floyd JC Jr, Fajans SS, Conn JW, Knopf RF, Rull J: Insulin secretion in response to protein ingestion. *J Clin Invest* 45:1479-86, 1966
20. Fajans SS, Floyd JC Jr, Knopf RF, Conn JW: Effect of amino acids and protein on insulin secretion in men. *Recent Prog Horm Res* 23:617-20, 1967
21. Nuttall FQ, Gannon MC, Wald JL, Ahmed M: Plasma glucose, and insulin profiles in normal subjects ingesting diets of varying carbohydrate, fat, and protein content. *J Am Coll Nutr* 4:437-50, 1985
22. Jenkins DJA, Wolever TMS, Wong GS, Kenshole A, Josse RF, Thompson LU, Lam KY: Glycemic responses to foods: possible differences between insulin-dependent and noninsulin-dependent diabetics. *Am J Clin Nutr* 40:971-81, 1984
23. Collier G, O'Dea K: The effect of coingestion of fat on the glucose, insulin, and gastric inhibitory polypeptide responses to carbohydrate and protein. *Am J Clin Nutr* 37:941-44, 1983
24. Service FJ, Molnar GD, Rosevear JW, Ackerman E, Gatewood LC, Taylor WF: Mean amplitude of glycemic excursions, a measure of diabetic instability. *Diabetes* 19:644-55, 1970
25. Crapo PA, Reaven G, Olefsky J: Postprandial plasma-glucose and -insulin responses to different complex carbohydrates. *Diabetes* 26:1178-83, 1977
26. Pacini G, Finegood DT, Bergman RN: A minimal-model-based glucose clamp yielding insulin sensitivity independent of glycemia. *Diabetes* 31:432-41, 1982
27. Staub H: Untersuchungen über den Zuckerstoff Wechsel des Menschen: I Mitteilung. *Z Klin Med* 91:44-60, 1921
28. Traugott K: Über das Verhalten des Blutzucker Spiegels bei wiederholter und verschiedener Art enteraler Zuckerzufuhr und dessen Bedeutung für die Leberfunction. *Klin Wochenschr* 1:892-94, 1922
29. Abaira C, Lawrence AM: The Staub-Traugott phenomena. III. Effects of starvation. *Am J Clin Nutr* 31:213-21, 1978
30. Nuttall FQ, Mooradian AD, Gannon MC, Billington C, Krezowski P: Effect of protein ingestion on the glucose and insulin response to a standardized oral glucose load. *Diabetes Care* 7:465-70, 1984
31. Pileggi VJ, Szustkiewicz CP: Carbohydrates. In *Clinical Chemistry, Principles and Technics*. 2nd ed. Henry RJ, Gannon DC, Winkelmann JW, Eds. New York, Harper & Row, 1974, p. 1284-325
32. Wursch P, DelVedovo S, Koellreutter B: Cell structure and starch nature as key determinants of the digestion rate of starch in legume. *Am J Clin Nutr* 43:25-29, 1986
33. Rosenthal SM, Ziegler EE: Effect of uncooked starches on the blood sugar of normal and of diabetic subjects. *Arch Intern Med* 44:344-50, 1929
34. Lii C-Y, Change S-M, Young YA-L: Investigation of the physical and chemical properties of banana starches. *J Food Sci* 47:1493-97, 1982
35. Gannon MC, Neil BJ, Westphal SA, Nuttall FQ: The insulin and glucose responses to meals of glucose + various proteins in type II diabetic subjects (Abstract). *Diabetes* 36 (Suppl. 1):12A, 1987